

# Surgical Treatment of Gastric Outlet Obstruction from A Large Gastroduodenal Trichobezoar: A Case Report

Mohammad Aslam, Mohd Atif Ansari, Nafees Ahamad\*, Sadiq Husain, Maikal Kujur

Department of General Surgery Jawaharlal Nehru Medical College and Hospital, Aligarh Muslim University, Aligarh, Uttar Pradesh, India 202002

\*Corresponding Author: Nafees Ahamad, Department of General Surgery Jawaharlal Nehru Medical College and Hospital, Aligarh Muslim University, Aligarh, Uttar Pradesh, India 202002

Received Date: September 08, 2022; Accepted Date: September 15, 2022; Published Date: September 23, 2022

**Citation:** Mohammad Aslam, Mohd Atif Ansari, Nafees Ahamad, Sadiq Husain, Maikal Kujur. (2022). Surgical treatment of gastric outlet obstruction from a large gastroduodenal trichobezoar: A casereport. *J. Clinical Oncology Case Reports*.1(2); DOI: 10.31579/cocr.2022/007

**Copyright:** © 2022, Nafees Ahamad, this is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

Bezoars are collection of foreign indigestible material accumulating in the gastrointestinal tract leading to intraluminal mass formation that impairs the gastrointestinal motility and can lead to obstruction of the small or the large bowel. These include phytobezoar, lactobezoar, pharmacobezoar, trichobezoar, and polybezoar. A trichobezoar is a mass of undigested hair within the gastrointestinal tract. Trichobezoars are seen in psychiatric young female with known to be trichophagic.


**Keywords:** gastroduodenal trichobezoar; phytobezoar; lactobezoar; pharmacobezoar; trichobezoar; polybezoar

## Introduction


Bezoars are collection of foreign indigestible material accumulating in the gastrointestinal tract leading to intraluminal mass formation that impairs the gastrointestinal motility and can lead to obstruction of the small or the large bowel. These include phytobezoar, lactobezoar, pharmacobezoar, trichobezoar, and polybezoar. A trichobezoar is a mass of undigested hair within the gastrointestinal tract. Trichobezoars are seen in psychiatric young female with known to be trichophagic [1]. Most common site of trichobezoar is the stomach. If trichophagia is not detected earlier then it can develop into Rapunzel Syndrome. Recurrence is overwhelming without psychiatric treatment after the surgery.

## Case presentation

A 21-year-old female patient known to be plucking and swallowing hairs for 1 year was admitted to the emergency department for nausea, vomiting, and history of weight loss. On physical examination, patient was having tachycardia and pallor. Per abdominal examination a large, hard mass in the epigastric and left hypochondrium. Complete blood count, electrolytes, kidney function tests, liver function tests, amylase and lipase were normal. Consequently, esophagogastroduodenoscopy (EGD) was done, revealing a collection of a huge hard hairball occupying the entire lumen of the stomach from the fundus through the pylorus reaching the duodenum. (Fig.1) On CECT a large mass measuring 30×13×12cm, well defined, multi-layered, inhomogenous, well circumscribed, solid appearing with pockets of air enmeshed within it, non-enhancing mass in the gastric lumen, extending upto proximal jejunum. The lesion was separated from the gastric walls by gastric fluid. No evidence of abnormal gastric mural thickening was noted (Fig. 1).



**Fig.1** Upper gastrointestinal Endoscopy demonstrating trichobezoar



**Fig.2** CECT Showing Huge Gastroduodenal Trichobezoar

Subsequently, midline laparotomy was planned. A 10 cm longitudinal gastrotomy was done on the anterior gastric wall, 8 cm from the pylorus. A large trichobezoar, filling the entire stomach and the duodenum and proximal part of jejunum was extracted (Fig. 3).



**Fig.3** *Gastroduodenal cast trichobezoar that was removed via laparotomy*

## Discussion

Trichobezoar is a rare disorder commonly occur in young adult females, with psychiatric problem. The site of hair pulling is most commonly from the scalp, but can occur other hairy areas of body area. (2,3) Trichobezoars in humans were first described from a post mortem by Swain in 1854. Formation in the stomach is that hair is undigestible and due to its smooth nature cannot be propelled with peristalsis. It leads to the accumulation of hair together with mucus and food, causing the formation of a trichobezoar [4,5]. In most cases, confined within the stomach. However, in some cases, it extends through the pylorus into the jejunum, ileum or even colon. This condition is called Rapunzel syndrome[6]. Affected patients remain asymptomatic for many years. Symptoms develop as the bezoar increases in size. The complications that have been reported over the years include gastric mucosal erosion, ulceration, and gastric outlet obstruction and perforation of the stomach or the small intestine, protein-losing enteropathy, and death. On clinical examination large mobile epigastric mass that may be indentable, the so-called Lamerton's sign. The most common diagnostic tool used in the is a CT scan, with a typical image showing a well-defined intraluminal ovoid heterogeneous mass with interspersed gas. Management options include endoscopic removal, laparoscopic removal (7), or via laparotomy. Conventional laparotomy is still the treatment of choice depending on the size and site of trichobezoar.

## Conclusion

Trichobezoars should be considered as a differential diagnosis in young females with abdominal pain and presence of an upper abdominal mass. Endoscopic or surgical removal can be performed safely and effectively. Psychiatric evaluation and therapy is required to prevent recurrence.

## References

1. S.Q. Falah, A. Amanullah, Huge trichobezoar in the stomach of a young female patient, *Gomal J. Med. Sci.* 14 (2016) 118-119.
2. S.M. Kinoo, B. Singh, Gastric trichobezoar: an enduring intrigue. *Case reports in gastrointestinal, Medicine* (2012).
3. I.S. Al-Janabi, M.A. Al-Sharbaty, M.M. Al-Sharbati, L.A. Al-Sharifi, A. (2004). Ouhtit, Unusual trichobezoar of the stomach and the intestine: a case report, *J. Med. Case Rep.*
4. S. Khattak, A. Kamal, Trichobezoar gomal, *J. Med. Sci.*
5. S. Mewa Kinoo, B. Singh. (2012). Gastric trichobezoar: an enduring intrigue, *Case Rep. Gastrointest. Med.*
6. Obinna Obinwa, David Cooper, Faraz Khan, James M. (2017). O'Riordan, Rapunzel syndrome is not just a mere surgical problem: a case report and review of current management, *World J. Clin. Cases* 5 (February (2)). 50-55,
7. R.R. Gorter, C.M.F. Kneepkens, E.C.J.L. Mattens, D.C. (2010). Aronson, H.A. Heij, Management of trichobezoar: case report and literature review, *Pediatr. Surg. Int.* 26 (May (5)). 457-463.

**Ready to submit your research? Choose ClinicSearch and benefit from:**

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

**At ClinicSearch, research is always in progress.**

Learn more <http://clinicsearchonline.org/journals/clinical--oncology-case-reports>



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.