

# The Role of Mutations on Gene Tcf4, In Pitt-Hopkins Syndrome

Shahin Asadi <sup>1\*</sup>, Shima Dolabi<sup>1</sup>, Mahtab Farrash Bashi Masjed<sup>1</sup>

Medical Genetics-Harvard University. Director of the Division of Medical Genetics and Molecular Optogenetic Research.

\*Correspondence Author: Shahin Asadi, Medical Genetics-Harvard University. Director of the Division of Medical Genetics and Molecular Optogenetic Research.

Received Date: September 30, 2022 | Accepted Date: October 15, 2022 | Published Date: October 28, 2022

Citation: Shahin Asadi, Shima Dolabi, Mahtab Farrash Bashi Masjed, (2022) The Role of Mutations on Gene Tcf4, In Pitt-Hopkins Syndrome. *International Journal of Clinical Research and Reports*.1(1); DOI:10.31579/2835-785X/004

Copyright: © 2022 Shahin Asadi, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

Pitt Hopkins Syndrome (PTHS) is a rare genetic and neurological disorder. Affected children have certain facial features, including mental retardation, developmental delays, impaired ability to speak, and recurrent seizures and respiratory pattern disorders. Other symptoms that may occur include poor coordination (ataxia), frequent non-functional hand movements, constipation, sleep disturbances, and severe myopia (myopia). Behavioral abnormalities are common, although children are often described as social and cheerful. Some children with autism spectrum disorders meet the criteria. The specific signs and symptoms of this disorder and their severity can vary from person to person. Pitt Hopkins syndrome is caused by a change (mutation) in the TCF4 gene. This mutation occurs spontaneously and is almost non-existent in a family. The disorder was first described in the medical literature in 1978, and its causative gene was discovered in 2008.

**Keywords:** pitt hopkins syndrome; tcf4 gene; ataxia, epilepsy; hyperventilation

## Introduction

Pitt Hopkins Syndrome is a genetic disease characterized by mental retardation and growth retardation, respiratory problems, recurrent seizures

(epilepsy), and distinctive facial features. People with Pitt Hopkins Syndrome have moderate to severe mental disabilities. Most people with mental retardation delay the development of mental and motor skills (psychomotor retardation).[1]



**Figure 1:** Picture of a boy with Pitt-Hopkins syndrome with a dysmorphic face. Clinical Signs and Symptoms of Pitt-Hopkins Syndrome

People with this syndrome have a delay in learning to walk and developing fine motor skills such as picking up small objects with their fingers. People with Pitt Hopkins syndrome are usually unable to speak. Some may learn to

say a few words. Many people with the disorder exhibit features of autism spectrum disorders that impair communication and social skills.[1]



**Figure 2:** Images of Pitt-Hopkins syndrome patients with related disorders.

Respiratory problems in people with Pitt Hopkins syndrome are characterized by periods of rapid breathing (hyperventilation) followed by periods of slow or stopped breathing (apnea). These periods can cause a lack of oxygen in the blood and lead to a bluish tinge to the skin or lips (cyanosis). In some cases, lack of oxygen can cause loss of consciousness. Some older people with Pitt Hopkins Syndrome have wide, round fingertips (clubfoot)

due to frequent periods of hypoxia. Respiratory problems only occur when a person is awake and usually first appear in mid-childhood, but can start as early as childhood. Excessive breathing and apnea can be caused by emotions such as excitement or anxiety or excessive fatigue (severe fatigue). [1,2]



**Figure 3:** Image of a man with Pitt-Hopkins syndrome with distinctive facial features.

Epilepsy also occurs in most people with Pitt Hopkins syndrome and usually begins in childhood, although it can be present from birth. People with Pitt Hopkins Syndrome have distinctive facial features that include thin eyebrows, sunken eyes, a prominent nose with a long nose bridge, a double raised curve on the upper lip (Cupid bow), a wide mouth with thick lips, and a long distance. The teeth are apart, and the ears are usually thick and cup-shaped. [1,2]

Children with Pitt Hopkins Syndrome typically have happy, irritating behaviors with frequent smiles, laughter, and touching gestures. However,


they can also experience anxiety and behavioral problems. Other features of Pitt Hopkins syndrome include constipation and other gastrointestinal problems, small head size (microcephaly), myopia (not seeing distant objects), eyes that are not visible in one direction (strabismus or lochia). Eyes), short stature and minor abnormalities of the brain. Affected people may also have small hands and feet, a crease in the palms of the hands, flat feet (pes planus), or unusual fleshy pads on the tips of the fingers and toes. Men with Pitt Hopkins syndrome may have cryptorchidism. [1,2]



**Figure 4:** Another view of patients with Pitt-Hopkins syndrome with related disorders. Etiology of Pitt Hopkins syndrome

A mutation in the TCF4 gene, located on the long arm of chromosome 18 at 18q21.2, causes Pitt Hopkins syndrome. This gene provides the instructions for the synthesis of a protein that binds to other proteins and then binds to specific regions of DNA to help control the activity of many other genes.


Based on DNA binding and gene control activities, the TCF4 protein is recognized as a transcription factor. The TCF4 protein is involved in the maturation of cells to perform specific functions (cell differentiation) and the destruction of cells themselves (apoptosis).[1,3]



**Figure 5:** Schematic of chromosome 18 where the TCF4 gene is located in the long arm of this chromosome as 18q21.2.

Mutations in the TCF4 gene impair the ability of proteins to bind to DNA and control the activity of specific genes. These disorders, especially the inability of the TCF4 protein to control the activity of genes involved in the development and function of the nervous system, contribute to the signs and symptoms of Pitt Hopkins syndrome. In addition, additional proteins interact with the TCF4 protein to perform specific functions. When the TCF4 protein is inactive, these other proteins are no longer able to function normally. It is also possible that the loss of normal proteins bound to non-functional TCF4

proteins may contribute to the characteristics of these conditions. Loss of a protein, particularly a protein synthesized by the ASCL1 gene, located on the long arm of chromosome 12 at 12q23.2, is thought to be associated with respiratory problems in people with Pitt Hopkins syndrome.[1,3] Pitt Hopkins syndrome follows an autosomal dominant inheritance pattern. Therefore, a copy of the TCF4 mutant gene (either parent) is required to cause the syndrome, and the chance of having a child with the autosomal dominant syndrome is 50% for each possible pregnancy.[1,3]




**Figure 6:** Schematic of the molecular and biochemical mechanism of the TCF4 gene in various organs of the body.

**Frequency of Pitt Hopkins Syndrome**

Pitt Hopkins syndrome is thought to be a very rare genetic disease. So far, about 500 people with this syndrome have been reported in the medical literature from around the world [1,4].

**Diagnosis of Pitt Hopkins Syndrome**

Pitt Hopkins syndrome can be diagnosed based on the clinical and physical findings of patients and some pathological tests. The most accurate way to diagnose this syndrome is a molecular genetic test for the TCF4 gene to check for possible mutations. [1,4].



**Figure 7:** Schematic of the predominant autosomal inherited pattern that Pete-Hopkins syndrome follows.

### Treatments for Pitt Hopkins Syndrome

The treatment and management strategy for Pitt Hopkins syndrome is symptomatic and supportive. Treatment may be performed with the efforts and coordination of a team of specialists including a pulmonologist, cardiologist, neurologist, oral and maxillofacial specialist, surgeons, and other health care professionals. There is no definitive treatment for this syndrome and all clinical measures are taken to alleviate the suffering of the patients. Genetic counseling is also essential for all parents who want a healthy baby. [1,4].

### Discussion and Conclusion

Pitt Hopkins Syndrome is a genetic disease characterized by mental retardation and growth retardation, respiratory problems, recurrent seizures (epilepsy), and distinctive facial features. People with Pitt Hopkins Syndrome have moderate to severe mental disabilities. Pitt-Hopkins syndrome is caused by an unexpected change (mutation) in the TCF4 gene. [1,5]. Genes provide instructions for creating proteins that play a critical role in many functions of the body. When a mutation of a gene occurs, the protein product may be faulty, inefficient, absent or overproduced. Depending upon the functions of the particular protein, this can affect many organ systems of the body, including the brain. The TCF4 gene creates a protein that is a transcription factor. This protein has an important role in various developmental processes of the body. It is highly expressed early during human development and is found throughout the central nervous system. The treatment and management strategy for Pitt Hopkins syndrome is symptomatic and supportive. Treatment may be performed with the efforts and coordination of a team of specialists including a pulmonologist,

cardiologist, neurologist, oral and maxillofacial specialist, surgeons, and other health care professionals. [1,6].

### References

1. Asadi S, (2022) Pathology in Medical Genetics Book, Vol 20, Amidi Publications, Iran
2. Goodspeed K, Newsom C, Morris MA, et al. (2018) Pitt-Hopkins syndrome: a review of current literature, clinical approach, and 23-patient case series. *J Child Neurol.* 33:233-244.
3. Bedeschi MF, Marangi G, Calvello MR, et al. (2018) Impairment of different protein domains causes variable clinical presentation within Pitt-Hopkins syndrome and suggests intragenic molecular syndromology of TCF4. *Eur J Med Genet.* 60:565-571.
4. de Winter CF, Baas M, Bijlsma EK, et al. (2016) Phenotype and natural history in 101 individuals with Pitt-Hopkins syndrome through an internet questionnaire system. *Orphanet J Rare Dis.* 11:37.
5. Rannals MD, Page SC, Campbell MN, et al. (2016) Neurodevelopmental models of transcription factor 4 deficiency converge on a common ion channel as a potential therapeutic target for Pitt-Hopkins syndrome. *Rare Dis.* 4:e1220468.
6. Marangi G, Zollino M. (2015) Pitt-Hopkins syndrome and differential diagnosis: a molecular and clinical challenge. *J Pediatr Genet.* 4:168-176.

**Ready to submit your research? Choose ClinicSearch and benefit from:**

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

**At ClinicSearch, research is always in progress.**

Learn more <https://clinicsearchonline.org/journals/international-journal-of-clinical-research-and-reports>



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.