

# Development of a Novel Beverage with Cactus Juice (*Opuntia ficus indica*) and Date Palm Sap (*Phoenix dactylifera L.*)

Ennouri M.<sup>1\*</sup>, Ennouri K.<sup>2</sup> and Triki M.A.<sup>1</sup>

<sup>1</sup> Olive tree institute, LR16IO01, University of Sfax, Tunisia

<sup>2</sup> Digital Research Center of Sfax, Technopark of Sfax, University of Sfax, Tunisia.

\*Corresponding Author: Ennouri M, Olive tree institute, LR16IO01, University of Sfax, Tunisia.

Received date: January 27, 2023; Accepted date: February 13, 2023; Published date: February 20, 2023

**Citation:** Ennouri M., Ennouri K and Triki M.A., (2023), Development of a Novel Beverage with Cactus Juice (*Opuntia ficus indica*) and Date Palm Sap (*Phoenix dactylifera L.*), *Clinics in Nursing*, 2(1) DOI:10.31579/ 2835-8147/013

**Copyright:** © 2023, Ennouri M. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

This study aimed to evaluate the possibility to produce a beverage combining *Opuntia ficus indica* juice (OFIJ) and date palm sap (*Phoenix dactylifera L.*) (DPS). Beverages (B1, B2, B3) were formulated by mixing the juice and date palm sap at different ratios (40:60%, 30:70%, 20:80%) whereas OFIJ and DPS were considered as the controls. Sensory evaluation using 5-point hedonic scale was performed and showed that the control DPS was the most appreciated in term of aftertaste and overall acceptability. Among the formulated beverages B3 is more appreciated than B2, followed by B1. Physico-chemical analysis showed that the sedimentation and serum separation increase considerably after 28 days of cold storage traducing the instability of beverages. Microbial analysis carried out during 42 days of cold storage revealed the absence of total coliforms, *Staphylococcus* and *Escherichia coli* for all drinks and that after 28 days of storage the beverages cannot be consumed.

**Keywords:** opuntia ficus indica; phoenix dactylifera l; juice; microbial count; sensory properties; stability

## 1.Introduction

Beverages based on juices are more and more consumed in regard to their healthy, tasty and refreshing character. Indeed, the average volume of juice consumed per person in 2023 is expected to amount of 4.66L. The volume growth of juices segment is expected to reach 1.8% in 2024 and it is expected to reach amount to 37.633ML by 2027 [1]. The most consumed juices, in volume, are the mixtures and smoothies, orange juice, apple juice, and in equal amounts grapefruit juice, pineapple juice and grape juice [1]. *Opuntia ficus indica* (OFI) presents high nutritional value, social, economic and environment benefits [2]. OFI can be used to stabilize soil and counter soil erosion, it can be used as a forage substitute, and it is considered as a dietary supplements and nutraceutical [3, 4]. *Opuntia ficus indica* juice (OFIJ) is rich in minerals, proteins, and in antioxidants such as polyphenols and flavonoids like reported in the work of Baccouche et al. [5] which evaluated the physical stability of whey-based *opuntia ficus indica* beverages. The study of Aloui et al [6] concerning twenty-two healthy men athletes, reporting the effect of OFIJ supplementation on oxidative stress, cardiovascular parameters and biochemical markers, concluded that supplementation with OFIJ decreases muscle damage caused by endurance exercise and reduces total and LDL-cholesterol. In the same vein, the more recent study of Bellafiore et al [7] has demonstrated that supplementation with OFIJ to women decreased the oxidative stress induced by intense exercise and improved balance in physically active women

Date palm sap (DPS) familiar in Tunisian population and called “legmi” is usually collected by traditional tapping technique, it is consumed in its fresh form as juice or as an alcoholic drink after auto fermentation [8]. DPS is a good source of antioxidant and possess antimicrobial activities particularly against Gram-positive bacteria and possess cytotoxic properties against HeLa human cell line [9]. Abdennabi et al [10] have isolated endophytic fungi from date palm sap and reported antifungal activities. More recently Ben Atitallah et al [11] reported the good potentiality of date palm sap for producing bioethanol.

The combination of *Opuntia ficus indica* juice and palm sap is so evident, since *Opuntia ficus indica* and *Phoenix dactylifera L.* are typical of semi-arid to arid regions and are adapted to growth in severe environment conditions. Each plant offers a juice which is very appreciated by the local population. When consumed separately the two juices are unstable, that's why we attempt to study the stability of their mixture which has never been reported in the literature.

The aim of our study is on the one hand to test the development of a new natural drink of high nutritional quality and to analyse its organoleptic properties, on the other hand to evaluate physico-chemical, and microbiological stability of drinks during storage at 5°C during 42 days.

## 2. Materials and methods:

### 2.1 Beverages:

Ripe cactus fruits (*Opuntia ficus indica*) were bought on the markets of sfax / Tunisia in early August and processed into juice. All the selected fruits are washed with water containing 0.01% chlorine, peeled manually, crushed, the mixed pulp was then passed through a strainer to retain the seeds. The extracted juice is filtered through cheesecloth and stored in the dark at -25°C until use. Exuded sap was obtained by a traditional tapping technique from palm grove (*Phoenix doctylifera L.*) and filtered through cheesecloth. The two juices were mixed in different percentage like reported in Table 1 and the beverages were formulation according to the flow chart presented in Figure 1. OFIJ and DPS were considered as controls. The photos of studied beverages were presented in Figure 2.

### 2.2 Physico-chemical analysis

The physico-chemical analysis concerned the determination of pH, turbidity, total soluble solids, sedimentation, and serum separation. The pH was determined by a pH meter (Eutech pH700) at 20°C [12], a turbidimeter (Aqua lytic AL450T-IR) was used for turbidity determination. Total soluble solids were determined at 20°C with a refractometer (ATAGO) and expressed as °Brix.

The sedimentation is carried out by centrifugation of the beverages at 4500 rpm for 20 min with a centrifuge (Universal 320R). The sedimentation corresponds to the pellet, its weight was calculated after elimination of the supernatant.

To evaluate serum separation, samples were placed in 10 ml plastic pipettes sealed at both ends each containing a beverage sample, pipettes were incubated at 5°C to assess serum separation by gravity and were inspected at various times during the 42 days of storage. When sedimentation takes place, a clear zone is observed in the upper part which is the sign of the instability of the drink [13]. All analyses are performed in triplicate.

### 2.3 Microbiological analysis

Beverage samples were aseptically removed from the bottles and 10 g from each sample were homogenized in 90 ml of sterile distilled water. Serial dilutions were prepared and 1 ml aliquots were plated in each specific medium and incubated at different temperatures, times and growth media as shown in Table 2.

### 2.4 Sensory analysis

The panel consisted of 25 people accustomed to consume fruit juices and cocktails. The panelists were asked to assess, by scoring on a scale of five, the intensity of seven descriptors; color, taste, smell, sweetness, texture, aftertaste and overall acceptability. During the sensory evaluation each panelist is separated from the others. When presented to panelists, samples are coded with 3-digit codes.

## 3. Results and discussion:

### 3.1 Sensory analysis

The results presented in Table 3 show that DSP is the more appreciated in term of aftertaste and overall acceptability respectively 4.21 and 4.55, whereas the OFIJ obtained the best note for odor. Regarding the formulated beverages, B3 has best notes for color (4.12), sweetness (4.68), texture (4.20) and overall acceptability (4.28). This result is explained by the fact that B3 contains 80% of DSP which was well appreciated by panelists, indeed DSP is appreciated by local population and so, it is recognised when analysed by panelists. We can conclude from the sensory evaluation that B3 is the best formulation, but it is important to evaluate the physical and microbial stability of this beverage during cold storage.

### 3.2 Physico-chemical beverages stability:

The evaluation of the physico-chemical characteristics of beverages during storage is presented in Table 4. The initial pH of OFIJ was 6.21 showing a

slight decrease with storage time reaching 6.03 after 42 days of storage. The pH of DSP is the more affected by storage, exhibiting a pronounced decrease (pH of 6.80 at day 1 vs. 4.21 at day 42). Regarding formulated beverages, the more affected by storage was B3 containing the highest DPS concentration and the less affected was B2.

The turbidity is influenced not only by storage time but also by the composition of the beverages, the more the percentage of OFIJ in beverage increases and the percentage of DSP decreases, the more the turbidity increases (Table 4). Over time, the turbidity increases more for B1 than for B2 and B3. It can be seen that the soluble solids content decreases for all the beverages during cold storage which is in accordance with the works of Ferreira et al [4] and Quiero et al [14] and can be attributed to the growth of microbial activity consuming thus the sugars.

Sedimentation and serum separation traduce the physical stability of beverages, these two parameters are very important for consumers and are determinant for purchasing intention. Sedimentation and serum separation of formulated beverages have varied respectively from 2.51 to 35.12% and from 14 to 16% for B3 during storage. The observed sedimentation increase is probably due to the complexation between the DPS components and those of OFIJ like reported by Baccouche et al [5] for whey and OFIJ.

### 3.3 Microbiological analysis:

The microbiological analysis was carried out on the three formulated beverages and the two controls (DSP and OFIJ) stored for a period of 42 days at 5°C. The microbiological analysis of the different samples showed that total coliforms, *Staphylococcus* and *Escherichia coli* are absent in all beverages.

During storage, the total mesophilic flora and the lactic acid bacteria increase especially during the last 14 days (Table 5). For this, we can say that the presence of these two floras in the drinks after 42 days can lead to the appearance of more or less important enzymatic activities in our product, hence the possibility of its alteration.

During the first thirteen days, the number of yeasts and moulds does not exceed 10 CFU in all drinks. This indicates stability probably due to the addition of potassium sorbate. However, between 28 and 42 days, the number of yeasts and moulds exceeded the standard which is 10<sup>2</sup> CFU/mL for pasteurized juices. We therefore conclude that from a microbiological safety point of view, that drinks have a shelf life of less than 28 days.

## Conclusion

The aim of our work is the elaboration of a new cocktail from two natural resources OFIJ and DPS that are found in our country but are badly exploited because of their rapid alteration. The sensory evaluation has demonstrated that B3, containing 20% *Opuntia ficus indica* juice and 80% date palm sap, is the more appreciated. Physico-chemical analysis showed that after 28 days of cold storage, sedimentation and serum separation increase considerably traducing the decrease of the beverage's stability. From a microbiological point of view, the analysis carried out on the beverages show that the products are safety until 28 days of storage.

## References:

1. Statista, 2023. <http://statista.com/outlook/cmo/non-alcoholic-drinks/juices/worldwide> consulted the 14/02/2023.
2. Nefzaoui, A. 2018, *Opuntia ficus-indica* productivity and ecosystem services in arid areas Review n. 36 – Italus Hortus 25 (3), 2018: 29-39. doi:10.26353/j.itahort/2018.3.2939
3. Jimenez-Aguilar, D.M.; Lopez-Martinez, J.M.; Hernandez-Brenes, C.; Gutierrez-Urbe, J.A.; Welti-Chanes, J. 2015. Dietary fiber, phytochemical composition and antioxidant activity of Mexican commercial varieties of cactus pear. Journal of food composition and analysis, 41, 66-73.
4. Ferreira, R.M.; Amaral, R.A.; Silva, A.M.S.; Cardoso, S.M.; Saraiva, J.A. (2022). Effect of high-pressure and thermal pasteurization on microbial and physico-chemical properties of

- Opuntia ficus-indica* juices. *Beverages* 8(4):84. <https://doi.org/10.3390/beverages8040084>.
5. Baccouche A, Ennouri M, Felfoul I, Attia H. (2013). A physical stability study of whey-based prickly pear beverages. *Food Hydrocolloids* 33:234–244. doi: 10.1016/j.foodhyd.2013.03.007.
  6. Aloui K, Abdelmalek S, Chtourou H, Souissi N. (2018). The effect of *Opuntia ficus-indica* juice supplementation on oxidative stress, cardiovascular parameters, and biochemical markers following yo-yo Intermittent recovery test. *Food Science and Nutrition*, 6:259-268. <https://doi.org/10.1002/fsn3.529>.
  7. Bellafiore, M., Pintaudi, A.M., Thomas, E. et al. (2021). Redox and autonomic responses to acute exercise-post recovery following *Opuntia ficus-indica* juice intake in physically active women. *Journal of the International Society of Sports Nutrition*, 18, 43. <https://doi.org/10.1186/s12970-021-00444-2>
  8. Makhoulf-gafsi, I., Mokni-ghribi, A., Bchir, B. et al. (2015). Foamability and Foam Stability of Male and Female Date Palm Sap (*Phoenix dactylifera L.*) During the Collection Period. *Food Biophysics* 10, 360–367. <https://doi.org/10.1007/s11483-015-9400-9>
  9. Makhoulf-Gafsi, I.; Baklouti, S.; Mokni, A.; Danthine, S.; Attia, H.; Blecker, C.; Besbes, S.; Masmoudi, M. (2016). Effect of ultrafiltration process on physico-chemical, rheological, microstructure and thermal properties of syrups from male and female date palm saps, *Food Chemistry*, 203, 175-182, <https://doi.org/10.1016/j.foodchem.2016.02.055>.
  10. Abdennabi, R., Triki, M.A., Ben Salah, R., Gharsallah, N. (2017). Antifungal activity of endophytic fungi isolated from date palm sap (*Phoenix dactylifera L.*). *EC Microbiology* 13.4: 123-131.
  11. Ben Atitallah, I., Arous, F., Louati, I., Zouari-Mechichi, H., Brysch-Herzberg, M., Woodward, S., Mechichi, T. (2021). Efficient bioethanol production from date palm (*Phoenix dactylifera L.*) sap by a newly isolated *Saccharomyces cerevisiae* X19G2. *Process Biochemistry*, 105, 102-112.
  12. AFNOR. (1970a). Produits dérivés des fruits et légumes : mesure électrométrique du pH
  13. avec l'électrode de verre, NF T 90e008, Ed. Paris: AFNOR.
  14. Koffi, E., Shewfelt, R., & Wicker, L. (2005). Storage stability and sensory analysis of UHT processed whey-banana beverages. *Journal of Food Quality*, 28, 386e401.
  15. Quieros, R.P.; Rainho, D.; Santos, M.D.; Fidalgo, L.G.; Delgadillo, I.; Saraiva, J.A. (2015). High pressure and thermal pasteurization effects on sweet cherry juice microbiological stability and physico-chemical properties. *High pressure research*.35, 69-77.

#### Ready to submit your research? Choose ClinicSearch and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At ClinicSearch, research is always in progress.

Learn more <https://clinicsearchonline.org/journals/clinics-in-nursing>



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.