

Iron Deficiency Among a Group Syrian Woman: Causes and Treatment

Sarah Sukkariyeh, Caren Latouf, Rim Harfouch*

Department of biochemistry and microbiology, college of pharmacy, Alsham private university (ASPU), Latakia, Syria.

***Corresponding Author:** Rim Harfouch, Department of biochemistry and microbiology, college of pharmacy, Alsham private university (ASPU), Latakia, Syria.

Received date: January 30, 2023; Accepted date: February 03, 2023; Published date: February 07, 2023

Citation: Sarah Sukkariyeh, Caren Latouf, Rim Harfouch, (2023). Iron Deficiency Among a Group Syrian Woman: Causes and Treatment. Journal of Skeleton System. 2(1). DOI: 10.58489/2836-2284/014

Copyright: © 2023 Rim Harfouch, This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Iron deficiency is one of the most common nutrient deficiencies in the world. The aim of our research was to identify the most common causes of iron deficiency anemia and the associated factors of lifestyle, dietary habits and menstrual cycle.

We selected a random sample of women attending the gynecology department at Tishreen University Hospital, and obtained the results of a complete blood count (HCT and HB) CBC for 100 participating women from April 2022 until the end of June 2022. As a result, 67% of women nearly two-thirds were anemic which is considered high for a sample randomly taken from the gynecological clinic in Tishreen university hospital.

Key words: Iron deficiency anemia; Syrian women; causes

Introduction

Iron deficiency is one of the most common nutrient deficiencies in the world. It can develop into anemia or remain iron deficiency. Hemoglobin concentration and/or erythrocyte volume decreases in iron- deficiency anemia due to insufficient iron availability to produce red blood cells [1].

Iron-deficiency anemia can arise from causes related to chronic blood loss, reduced iron intake or absorption, increased iron demand, or chronic inflammation. Iron-deficiency anemia also occurs in patients with chronic, inflammatory, cancer, or renal disease because inflammatory processes impair the delivery of iron to red cell precursors, regardless of iron stores [2].

Females are particularly vulnerable to iron deficiency and can suffer serious lifelong health consequences if not diagnosed and treated appropriately.

Perimenopausal women presenting with anemia, the most common cause reported is heavy menstrual blood loss. Menstrual disorder accounts for 5%–10% of the women presenting with iron deficiency anemia (IDA) in the perimenopausal age group [3].

In developing countries like India where the nutritional Iron deficiency is very much prevalent in the perimenopausal age group women, menorrhagia acts as

an aggravating factor leading to severe anemia. Anemia contributes to increase morbidity and mortality in this age group.

Materials and Methods

The aim of our research was to identify the most common causes of iron deficiency anemia and the associated factors of lifestyle, dietary habits and menstrual cycle. We selected a random sample of women attending the Women's Department at Tishreen University Hospital, and obtained the results of a complete blood count (HCT and HB) CBC for 100 participating women from April 2022 until the end of June 2022.

The survey was conducted under the medical ethics applied in Tishreen university hospital and after obtaining the patients consent [5].

Results

Distribution of the study sample according to the hemoglobin concentration The study sample was divided into 3 categories: higher than <10g>12g\dl, 10-12g\dl.

According to the WHO criteria for assessing iron- deficiency anemia (less

than (12 g/dl)), it was found that the proportion of normal women without iron- deficiency anemia is 33% (hemoglobin above 12g/dl), while the percentage of women with iron- deficiency anemia is 67% which is two-thirds

which is considered high for a sample randomly taken from the gynecological clinic in Tishreen University Hospital




Figure 1: represents the distribution of the study sample according to the concentration of hemoglobin

2. Distribution of the study sample according to family status

The study sample was divided into 3 categories:

single, married, not pregnant and married, pregnant.

The percentage of single women was 50%, of them 24 had iron deficiency

anemia. Surprisingly, 43 out of 50 married women were affected by iron deficiency anemia in our study, and we noticed that 9 pregnant women with iron deficiency anemia out of 13 pregnant women, i.e. 69%, and this percentage calls for emphasizing the necessity of Pregnant women take iron supplements during pregnancy due to the increased need to secure the requirements of the fetus.




Figure 2: Represents the distribution of the study sample according to family status

Distribution of the study sample according to the abundance of the menstrual cycle

According to the number of days of the menstrual cycle:

The study sample was divided into two groups according to the duration of the menstrual cycle:

3-5 days and 6-8 days.

It was found that 15 patients with iron deficiency anemia out of 35 women have a period of 3-5 days, or 42.8%, and 51 patients with iron deficiency anemia out of 64 women whose cycle lasts 6-8 days (80%). This may indicate that the length of the menstrual period is associated with the possibility of iron deficiency anemia, and there was in our study one patient in the menopausal stage (amenorrhea) with uterine fibrosis associated with abnormal bleeding, which explains her iron deficiency anemia.




Figure 3: represents the distribution of the study sample according to the number of days of the menstrual cycle

4. Distribution of study sample according to the intensity of the flow

The patients were asked about the intensity of their menstrual flow and were divided into categories:

light 7%, normal 39%, heavy 41%, very heavy 12% in addition to one patient in menopause.

In comparison, 3 out of 7 light-flowing women had iron deficiency anemia (with hemoglobin of 10-12 g/dl).

While those with heavy menstrual cycles, 30 out of 41 suffer from iron deficiency anemia (73%), while those with very heavy menstrual cycles, 12 out of 12 suffer from iron deficiency anemia, knowing that 7 of them have hemoglobin less than 10g/dl, which is a logical result as Heavy bleeding will lead to the inability of the bone marrow to compensation red blood cells.




Figure 4: Represents the distribution of the study sample according to the abundance of menstrual flow

Distribution of the study sample according to the intake of antacids (Proton pump inhibitors)

The study sample was divided into two groups according to the intake of antacids, and the percentage of those who are taking antacids was 15%, 12 of them were patients with iron deficiency anemia (80%) and it is considered high. It was found through a study that there is a relationship between taking

antacids and ID, as the cause Iron deficiency is a decrease in iron absorption from the gastrointestinal tract with a decrease in gastric acid secretion due to the use of PPIs. As for the percentage of women who do not take antacids, 85% of them were 55 patients suffering from iron deficiency anemia (64%).




Figure 5: Sample distribution of the study by taking antacids

6. Distribution of the study sample according to diet (meat intake)

The study sample was divided into two groups according to meat consumption, and the percentage of those who ate meat was 69%, of them 42 patients with iron deficiency anemia (i.e., 60%), while the percentage of those who did not eat meat was 31%, of them 25 were patients with iron deficiency anemia (i.e., 80%) is considered high.

This diet needs to be discussed, as it should be emphasized that vegetarians should replace meat with iron supplements or replace with iron-rich legumes.

It was noted among the participants that 4 of them suffer from malabsorption (celiac disease) and 3 of them suffer from iron deficiency anemia, this

condition does not benefit from diet modification and iron supplements must be given.

7. Distribution of the study sample according to treatment

It was found previously that the percentage of women with iron-deficiency anemia was 67%, 36 patients of them were treated (i.e. 54%), 4 patients of them were treated intravenously (their hemoglobin concentration was less than 10g/dl) and it is considered an emergency case that requires raising iron stores quickly (oral iron needs 4 weeks to take effect), while the number of untreated patients was 31 (46%), and this lead to the need to spread awareness among these patients of the necessity to take iron compounds.




Figure 6: represents the distribution of the study sample according to meat intake




Figure 7: represents the distribution of the study sample according to treatment

8. Distribution of the study sample according to the side effects of the treatment

Orally treated patients were divided into 4 categories according to side effects:




Figure 8: represents the distribution of the study sample according to the side effects of treatment

constipation and/or indigestion, diarrhea, abdominal pain, no symptoms.

The number of women suffering from constipation and/or indigestion was 16 patients (47%), the number of women suffering from diarrhea was 6 patients (18%), who suffered from abdominal pain were 3 patients (9%), and those who did not complain of any symptoms were 9 patients (26%). Oral iron supplements are known to cause constipation and indigestion in high rates in patients.

Conclusion

Heavy menstrual bleeding is one of the most important factors contributing to iron deficiency anemia. Measures are needed to alleviate menstrual disorders, and improve iron status. Oral contraceptives can be part of a strategy to reduce anaemia, particularly for adolescents at high risk of unwanted pregnancies.

References

1. Todd, T., & Caroe, T. (2007). Newly diagnosed iron deficiency anaemia in a premenopausal woman. *BMj*, 334(7587), 259-259.
2. Lyseng-Williamson, K. A., & Keating, G. M. (2009). Ferric carboxymaltose: a review of its use in iron-deficiency anaemia. *Drugs*, 69, 739-756.
3. Mishra, V., Verneker, R., Gandhi, K., Choudhary, S., & Lamba, S. (2018). Iron deficiency anemia with menorrhagia: Ferric carboxymaltose a safer alternative to blood transfusion. *Journal of mid- life health*, 9(2), 92.
4. Mishra V, Verneker R, Gandhi K, Choudhary S, Lamba S. Iron deficiency anemia with menorrhagia: Ferric carboxymaltose a safer alternative to blood transfusion. *J Mid-life Health* 2018; 9:92-6.
5. Harfouch, R. M. (2021). Polycystic ovary syndrome: Strategies of treatments. *Open J Clin Med Images*, 1(1), 1001

Ready to submit your research? Choose ClinicSearch and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At ClinicSearch, research is always in progress.

Learn more <http://clinicsearchonline.org/journals/journal-of-clinical-anatomy>



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.