

# Antiarrhythmic Drugs vs. Ablation for Atrial Fibrillation

Ghazala Irfan

National Institute of Cardiovascular Diseases (NICVD), Karachi, Pakistan.

\*Corresponding Author: Ghazala Irfan, National Institute of Cardiovascular Diseases (NICVD), Karachi, Pakistan.

Received date: July 29, 2025; Accepted date: August 19, 2025; Published date: August 29, 2025

Citation: Ghazala Irfan, (2025), Antiarrhythmic drugs vs. Ablation for atrial fibrillation, *Journal of Heart and Vasculature*, 4(4); DOI:10.31579/2834-8788/33

Copyright: © 2025, Ghazala Irfan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

Atrial fibrillation (AF) is a common and potentially life-threatening cardiac arrhythmia that affects millions of individuals worldwide. The treatment landscape for AF has traditionally comprised antiarrhythmic drugs (AADs), which aim to restore and maintain normal heart rhythm. However, over the past two decades, catheter ablation has emerged as an alternative therapeutic approach in the management of AF. This article systematically compares the effectiveness, safety, and long-term outcomes of AADs and catheter ablation as treatment strategies for atrial fibrillation.

**Keywords:** atrial fibrillation; antiarrhythmic drugs; catheter ablation

## Introduction

Atrial fibrillation (AF) is characterized by disorganized electrical activity in the atria, causing irregular and often rapid heart rhythm. Over the years, pharmacological therapy with antiarrhythmic drugs (AADs) has been the mainstay of AF treatment. Nonetheless, catheter ablation has gained significant momentum due to its ability to directly target the arrhythmogenic substrate within the heart.<sup>1</sup>

### Efficacy:

1. Antiarrhythmic Drugs: Various classes of AADs inhibit specific ion channels or receptors to normalize cardiac electrical conduction. While AADs can effectively restore sinus rhythm in many patients, their long-term success in maintaining sinus rhythm is modest, with recurrence rates ranging from 40% to 60% within one year.<sup>2</sup>
2. Catheter Ablation: Ablation procedures selectively target and destroy the abnormal electrical pathways contributing to AF. In well- selected patients, catheter ablation has demonstrated higher efficacy rates than AADs, particularly in paroxysmal AF, with success rates reaching 70-80% or more after a single procedure.<sup>2</sup>

### Safety Profile:

1. Antiarrhythmic Drugs: Although generally well- tolerated, AADs have notable adverse effects, including pro-arrhythmia, organ toxicity, and drug-drug interactions. These side effects often
2. limit their usage or require careful patient monitoring.<sup>3</sup>
- Catheter Ablation: While catheter ablation is generally safe, it has a finite risk of procedural complications, such as vascular injury, cardiac perforation, and stroke.

Nonetheless, the overall complication rate is low, ranging from 1% to 5%, and is further decreasing with advancements in technology and operator experience.<sup>3</sup>

3. Quality of Life and Symptom Control:
4. Antiarrhythmic Drugs: Effective rate control with AADs can significantly improve symptom burden and quality of life in patients with persistent or permanent AF.<sup>4</sup>
5. Catheter Ablation: Successful ablation procedures can offer long-term freedom from AF symptoms, eliminating or reducing the dependence on AADs and their associated side effects.<sup>4</sup>

### Long-Term Outcomes:

1. Antiarrhythmic Drugs: AAD therapy is often lifelong, and patients may require multiple medication adjustments due to loss of efficacy or intolerable side effects. However, AADs remain a treatment option for patients with contraindications for ablation or in those who prefer medical management.<sup>4</sup>
2. Catheter Ablation: Successful ablation procedures have been associated with excellent long-term outcomes, with sustained freedom from AF achieved in approximately 60-70% of patients at 1-5 years of follow-up. Repeat ablations may be necessary in some cases.<sup>4</sup>

## Conclusion:

While both antiarrhythmic drugs and catheter ablation have their advantages and disadvantages in the management of atrial fibrillation, catheter ablation has emerged as a promising treatment strategy, offering higher success rates and long-term symptom control compared to AADs. However, individual

patient characteristics, preferences, and comorbidities should guide the selection of the optimal treatment approach, ensuring personalized and comprehensive management of atrial fibrillation. Further research is needed to refine patient selection criteria, improve ablation techniques, and optimize the use of antiarrhythmic drugs to enhance treatment outcomes for AF patients.

## References

1. Asad ZUA, Yousif A, Khan MS, Al-Khatib SM, Stavrakis S. Catheter Ablation Versus Medical Therapy for Atrial Fibrillation: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Circ Arrhythm Electrophysiol.* 2019;12(9): e007414.
2. Andrade JG, Deyell MW, Macle L, Wells GA, Bennett M, Essebag V, et al. EARLY-AF Investigators. Progression of Atrial Fibrillation after Cryoablation or Drug Therapy. *N Engl J Med.* 2023;388(2):105-16.
3. Charitakis E, Metelli S, Karlsson LO, Antoniadis AP, Rizas KD, Liuba I, et al. Comparing efficacy and safety in catheter ablation strategies for atrial fibrillation: a network meta-analysis. *BMC Med.* 2022;20(1):1-3.
4. Deshpande R, AlKhadra Y, Singanallur P, Botchway A, Labedi M. Outcomes of catheter ablation versus antiarrhythmic therapy in patients with atrial fibrillation: a systematic review and meta- analysis. *J Interv Card Electrophysiol.* 2022;65(3):773-802.

### Ready to submit your research? Choose ClinicSearch and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

**At ClinicSearch, research is always in progress.**

Learn more <http://clinicsearchonline.org/journals/journal-of-clinical-anatomy>



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.